



TREE SPECIES (SCIENTIFIC NAME)

*Viburnum lehmannii*

COLOMBIAN COMMON NAME

Sauco de monte

TREE FAMILY

VIBURNACEAE

AVERAGE LEAF SIZE (CM)

10cm × 5cm  
Length Width

DISTRIBUTION



NATIVE TO COLOMBIA

ELEVATIONAL RANGE (M)

1100–3650m

TREE HEIGHT

MEDIUM (20–35M)



NATIVE TO

Region:

Colombia: Boyacá, Cauca, Huila, Valle del Cauca

COFFEE AGROFORESTRY INFORMATION

COFFEE SYSTEM



ARABICA

COFFEE IMPACT

Unknown

TREE MANAGEMENT

Propagated by seeds and cuttings.

CULTIVATION



NATURAL

TREE BENEFITS AND USES

FARMER USES



Livestock Forage, Lumber, Ornamental, Product

Bark dye is used for tanneries and fruit dye for handicrafts. Used for construction, carpentry and cabinetmaking.

FARM SERVICES



Windbreak, Carbon Capture, Water Protection

Used for water protection and decontamination

BIODIVERSITY BENEFITS

✓ YES

Attracts biological control agents, Attracts seed dispersers, Attracts pollinators. Visited by generalist and open-area birds looking for insects to feed on. Sparrows, thrushes and seedeaters are observed. Its flowers attract insects and bats. In addition, it is an endemic species to Colombia, which means that it is not found anywhere else in the world!

Last Updated: February 20, 2025

Plants of the World Online (POWO). (2024). Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet. Retrieved from <http://www.plantsoftheworldonline.org/>; International Union for Conservation of Nature (IUCN). (2024). IUCN Red List. Published on the Internet. Retrieved from <https://www.iucnredlist.org/>; World Flora Online. (2024). World Flora Online. Published on the Internet. Retrieved from <https://www.worldfloraonline.org/>; Global Biodiversity Information Facility (GBIF). (2024). GBIF. Published on the Internet. Retrieved from <https://www.gbif.org/es/>