



TREE SPECIES (SCIENTIFIC NAME)

*Nephelium lappaceum*

ENGLISH COMMON NAME

Rambutan

INDONESIAN COMMON NAME

Rambutan

TREE FAMILY

SAPINDACEAE

AVERAGE LEAF SIZE (CM)

6-18cm \* 4-7.5cm

Length Width

ELEVATIONAL RANGE (M)

0-600m

TREE HEIGHT

SMALL (10-20M)



DISTRIBUTION



NATIVE TO INDONESIA

NATIVE TO

Region: Asia

Malaysia, Singapore, Thailand

Indonesia: Borneo, Sumatra

EXOTIC IN

SE Asia: Laos, Myanmar, Philippines, Vietnam

Indonesia: Java, Lesser Sunda Islands, Sulawesi

## COFFEE AGROFORESTRY INFORMATION

COFFEE SYSTEM



ROBUSTA



ARABICA

COFFEE IMPACT



BENEFICIAL TO COFFEE

REGENERATION



REGENERATES NATURALLY

CULTIVATION



PLANTED

PREVALENCE



COMMON IN COFFEE AGROFORESTRY

TREE MANAGEMENT

Seedlings available from fruit/commodity nurseries. Seed propagation not recommended. Produces long, upright growth so early pruning/training is recommended to form an open center. After harvest, fruited twigs should be pruned back to stimulate new growth (up to 4 new side shoots), which allows 22% of the shoots to produce fruit in the following season. Remove dead branches and water suckers regularly. Mulch during establishment and dry periods. Do not mulch prior to flowering. Fertilizer recommended for young trees (200 g nitrogen, 25 g of phosphate, and 100 g potassium per tree per year). For the first 4 years, apply fertilizer every 3 months. For fruiting trees, apply fertilizer (200 g N, 25 g P, and 130 g K) per tree per year of age. A quarter of the yearly fertilizer should be applied 4 weeks after fruit set, half applied immediately after harvest, and the remaining quarter 9 weeks post-harvest. Additionally, 0.4 kg of dolomite per tree per year of age should be applied during slow growing months. Glyphosate herbicide should never be used near the drip line at any stage as it could cause a severe yellowing and abscission of the lower leaves. Fertilizer and dolomite use should constant after 10 years.

## TREE BENEFITS AND USES

FARMER USES

**Food, Fuelwood, Lumber, Medicinal, Ornamental, Product**

Fruit, oil, stimulant are used and consumed. Wood used in civil construction. Roots used to treat fever, the bark as an astringent for disease of the tongue, and the leaves as compresses for headache. Young shoots are used as a green dye for silk which has already been dyed yellow with turmeric (*Curcuma longa*). Used ornamentally for its fruits. Fruits can be sold at local markets and 40-80% of yearly production is generally marketed.

FARM SERVICES

**Coffee Shade**

BIODIVERSITY BENEFITS

Unknown

Last Updated: April 27, 2021

Image: Joanna Durant Digital Image © Board of Trustees, RBG Kew <http://creativecommons.org/licenses/by/3.0/>