



TREE SPECIES (SCIENTIFIC NAME)

Amburana cearensis

PERU COMMON NAME

Ishpingo

TREE FAMILY

FABACEAE

ELEVATIONAL RANGE (M)

350–500M

AVERAGE LEAF SIZE (CM)

Unknown

TREE HEIGHT

MEDIUM (20–35M)

DISTRIBUTION



NATIVE TO

Region: Americas**Latin America:** Argentina, Bolivia, Brazil, Paraguay, Peru

COFFEE AGROFORESTRY INFORMATION

COFFEE SYSTEM



COFFEE IMPACT

Unknown

CULTIVATION



PREVALENCE

Unknown

TREE MANAGEMENT

Plant the seeds in partial shade. An 80% germination rate can be expected for fresh seeds, with seeds sprouting within 15 to 25 days. Seedlings may grow slowly.

TREE BENEFITS AND USES

FARMER USES

**Lumber, Medicinal, Product**

It is commonly harvested from the wild for its attractive wood, which is used for luxury furniture, decorative veneer, construction, carving, joinery and to make boxes. Tree resin oil is used in the treatment of colds, coughs, bronchitis, asthma, and lung diseases. The seeds are used to treat asthma, relieve spasms, support cardiac function, and to stimulate menstrual flow. The seeds are used as perfume and the seeds soaked in water are used to kill mosquito larvae.

FARM SERVICES

Unknown

BIODIVERSITY BENEFITS

No

Last Updated: August 22, 2023

Image: Herbarium Catalogue Specimens Digital Image © Board of Trustees, RBG Kew <http://creativecommons.org/licenses/by/3.0/>

Jezeer, Rosalien. (2018). PhD dissertation: Shedding Light on Shade- Reconciling Livelihoods and Biodiversity in Coffee Agroforests. 10.13140/RG.2.2.28895.71844.;

Mendoza, A. L. M., & Guerrero, M. H. (2010). Catálogo de los árboles y afines de la Selva Central del Perú. Arnaldoa, 17, 203-242.;

Tropical Plants Database, Ken Fern. tropical.theferns.info/tropical.php?id=Amburana+cearensis