

TREE SPECIES (SCIENTIFIC NAME)

Acrocarpus fraxinifolius

ENGLISH COMMON NAME

The Pink Cedar

INDONESIAN COMMON NAME

Delimas, Madang pariek

TREE FAMILY

FABACEAE

AVERAGE LEAF SIZE (CM)

8.5cm × 30cm

Length Width

ELEVATIONAL RANGE (M)

0-1500 M

TREE HEIGHT

LARGE (> 35M)



DISTRIBUTION



EXOTIC IN INDONESIA

NATIVE TO

Region: Asia

Laos, Myanmar, Thailand, Vietnam

EXOTIC IN

Indonesia: Java, Sumatra

TREE MANAGEMENT

COFFEE AGROFORESTRY INFORMATION

COFFEE SYSTEM



COFFEE IMPACT

Unknown

REGENERATION

TEGENERATES NATURALLY



CULTIVATION



PREVALENCE



COMMON IN COFFEE AGROFORESTRY

TREE BENEFITS AND USES

FARMER USES







Livestock Forage, Fuelwood, Lumber

The wood is heavy, moderately hard, and compact with its whitish sapwood and bright red to brownish-red heartwood making it very decorative. The wood is not very durable and is prone to attack by fungi and insects, but it is easy to work with tools and is well suited for wood crafts, general construction, furniture, shingles, and beehive frames. The foliage can be used as livestock fodder.

FARM SERVICES











Coffee Shade, Windbreak, Soil Improvement, Erosion Control, Reforestation, Nitrogen Fixation

Regenerates naturally on burned or newly exposed soils. Periodic weeding

required until well established as saplings. Thinning should first be performed 3-4 years after initial planting then continuously until the stand is well developed as the tree requires a large crown for optimal growth. Competes

with other crops. Should not be planted near houses as dry branches drop off

Erosion control: recommended for reinforcing riverbanks and stabilizing terraces

Soil improvement: leaves are suitable for mulching

BIODIVERSITY BENEFITS

Unknown

Last Updated: April 26, 2021

 $Image: http://specimens.kew.org/herbarium/K000789066\ The\ Herbarium\ Catalogue,\ Royal\ Botanic\ Gardens,\ Kew.\ Published\ on\ the\ Internet\ http://www.kew.org/herbcat\ [accessed\ on\ 10]{Abstractions} and the second of t$ 12 2020]