



TREE SPECIES (SCIENTIFIC NAME)

# Flemingia macrophylla

ENGLISH COMMON NAME

Large leaf flemingia

INDONESIAN COMMON NAME

Hahapaan , Pok kepokan

TREE FAMILY

**FABACEAE**

AVERAGE LEAF SIZE (CM)

**7-11cm** × **3-6cm**  
Length Width

DISTRIBUTION



**NATIVE TO INDONESIA**

ELEVATIONAL RANGE (M)

**0-2000m**

TREE HEIGHT

**SHRUB (1-10M)**



NATIVE TO

**Region:** Asia

Cambodia, Laos, Malaysia, Myanmar, Papua New Guinea, Philippines, Singapore, Thailand, Vietnam

**Indonesia:** Borneo, Java, Lesser Sunda Islands, New Guinea, Sulawesi, Sumatra

## COFFEE AGROFORESTRY INFORMATION

COFFEE SYSTEM



**COFFEE**

COFFEE IMPACT



**BENEFICIAL TO COFFEE**

TREE MANAGEMENT

Propagated by seed. Scarification recommended before sowing to speed up and improve germination Pour a small amount of warm water (not boiling) on seeds and then soak them for 12 - 24 hours. Weed control required first 6 months. Mulching (at a rate of 3 t/ha) can effectively control weeds for ~3 months. Plants should not be cut at intervals less than 40 days but can be cut more frequently than every 3 months

REGENERATION

Unknown

CULTIVATION



**PLANTED**

PREVALENCE

Unknown

## TREE BENEFITS AND USES

FARMER USES



**Food, Medicinal**

Leaves are sometimes eaten. The leaves reduce fever and are used for treating postpartum fever as well as paralysis and pain in the joints. A decoction of the leaves is used to bathe sores and swellings and the entire plant is given to relieve stomachache

FARM SERVICES



**Coffee Shade, Windbreak, Soil Improvement, Erosion Control, Nitrogen Fixation, Weed Control**

*Soil improvement:* provides mulch, moisture conservation, reduces soil temperature

BIODIVERSITY BENEFITS

Unknown

Last Updated: April 27, 2021

Image: Dipankar Borah© Digital Image © Board of Trustees, RBG Kew <http://creativecommons.org/licenses/by/3.0/>