

#### TREE SPECIES (SCIENTIFIC NAME)

# Artocarpus altilis

## TREE FAMILY

MORACEAE

AVERAGE LEAF SIZE (CM) 42.2-78.6cm × 30.2-59.2cm Length Width

ELEVATIONAL RANGE (M)

0-1550м



## **COFFEE AGROFORESTRY INFORMATION**

COFFEE SYSTEM	COFFEE IMPACT
🌗 ROBUSTA 🌎 ARABICA	✓ BENEFICIAL TO COFFEE
REGENERATION	CULTIVATION
REGENERATES NATURALLY	🍹 PLANTED

PREVALENCE

Unknown

## **TREE BENEFITS AND USES**

#### FARMER USES

### 💥 🚌 🚳 🕂 🥡

#### Food, Livestock Forage, Lumber, Medicinal, Product

The fruit can be eaten raw, boiled, pickled, candied, marinated, steamed or roasted. The fruit is fried and made into sukun chips. The leaves are eaten by cattle, goats, pigs, horses, and elephants, and excess fruit waste can be fed to animals. Horses will eat the bark, young branches, and shoots, so they should be kept away from new plantings. The wood is very light, durable, and resistant to termites. Is used in house, furniture, and canoe construction. Toasted flowers are rubbed on the gums around aching teeth to ease pain, the latex and juice from crushed leaves are used to treat ear infections, skin ailments, and fungal diseases. The latex is massaged into the skin to treat broken bones, sprains, and back pain. The latex is diluted and taken internally to treat stomach problems. The root is an astringent used in skin ailments, and the bark treats headache.

Last Updated: April 26, 2021 Image: Ori Fragman-Sapir© Digital Image © Board of Trustees, RBG Kew http://creativecommons.org/licenses/by/3.0/

#### ENGLISH COMMON NAME

Breadfruit

INDONESIAN COMMON NAME

Sukun, Kelur, Timbul

#### DISTRIBUTION



#### NATIVE TO

Region: Asia Papua New Guinea, Philippines Indonesia: Lesser Sunda Islands, Maluku Islands, New Guinea, Sulawesi

EXOTIC IN

SE Asia: Laos, Malaysia, Singapore, Vietnam Indonesia: Java, Sumatra

#### TREE MANAGEMENT

Regardless of propagation method, young plants do best under shade, but require full sun once established. Land preparation consists of deep ploughing followed by harrowing. Should be planted at the beginning of the rainy season with optional supplemental irrigation during establishment. Spacing of approximately 100 trees/ha at 12 x 8 m or 10 x 10 m apart. Can be grown on a variety of soils, but thrives on alluvial, coastal, well-drained sandy loam, or clay loam soils. Good drainage is essential as trees may shed their fruit when the soil is excessively wet. Some cultivars (especially interspecific hybrids) have adapted to shallow, calcareous soils and appear to tolerate high saline conditions. After establishment they require little management. Insect pollinated.

#### FARM SERVICES



#### Coffee Shade, Soil Improvement

#### **BIODIVERSITY BENEFITS**

Unknown