

TREE SPECIES (SCIENTIFIC NAME)

Carallia brachiata

ENGLISH COMMON NAME

Corkybark

INDONESIAN COMMON NAME

Perepat Hutan, Ampana, Aroi, Aru, Bara, Belengkeching, Bepinang, Beranggae, Dabong, Ganggang putih, Kelai, Kitir-kitir, Massulie, Mundar, Mundar burung, Pacar, Petis, Rabong, Sawar bubu, Selunser, Tampilas, Tulang ular (Borneo), Manggisan

TREE FAMILY

RHIZOPHORACEAE

AVERAGE LEAF SIZE (CM)

4-10см × 4-6см

Length Width

ELEVATIONAL RANGE (M)

0-800 M

TREE HEIGHT

LARGE (> 35M)





NATIVE TO INDONESIA

NATIVE TO

DISTRIBUTION

Region: Asia

Cambodia, Laos, Malaysia, Myanmar, Papua New Guinea, Singapore, Thailand,

Vietnan

Indonesia: Borneo, Java, Lesser Sunda Islands, Maluku Islands, New Guinea,

Sulawesi, Sumatra

COFFEE AGROFORESTRY INFORMATION

COFFEE SYSTEM

ROBUSTA

COFFEE IMPACT

Unknown

CULTIVATION

TREE MANAGEMENT

Seeds dispersed by animals. Tree of late successional forest.

REGENERATION

TEGENERATES NATURALLY

PREVALENCE

Unknown

TREE BENEFITS AND USES

FARMER USES







Fuelwood, Lumber, Medicinal

The wood is used for furniture, floor paneling, and parquet flooring. Leaves used medicinally.

FARM SERVICES

Unknown

BIODIVERSITY BENEFITS

Unknown

Last Updated: April 27, 2021

Image: Helen Hewitt Digital Image @ Board of Trustees, RBG Kew http://creativecommons.org/licenses/by/3.0/