

TREE SPECIES (SCIENTIFIC NAME)

# Flacourtia rukam

AVERAGE LEAF SIZE (CM)

Width

12.25cm × 6cm

Length

## SALICACEAE

TREE FAMILY

ELEVATIONAL RANGE (M)

0-1500м



### **COFFEE AGROFORESTRY INFORMATION**

COFFEE SYSTEM	COFFEE IMPACT
🌗 ROBUSTA	Unknown
REGENERATION	CULTIVATION
C REGENERATES NATURALLY	J PLANTED

#### INDONESIAN COMMON NAME

Rukam, Ganda rukem, rukam (Java), Klang tatah kutang (Borneo)

#### DISTRIBUTION



#### NATIVE TO

Region: Asia Malaysia, Papua New Guinea, Philippines, Singapore Indonesia: Borneo, Java, Lesser Sunda Islands, Maluku Islands, New Guinea, Sulawesi, Sumatra

#### TREE MANAGEMENT

Propagated by seed and germinates readily and quickly. Shoot growth occurs in synchrony. Young plants bear much larger leaves than the full-grown tree which is often sparsely foliated. In Java the trees flower between June and August. Fruit ripens after about 14 weeks. Seeds dispersed by animals. Long-lived tree of late successional forest

#### PREVALENCE

Unknown

## **TREE BENEFITS AND USES**

#### FARMER USES

💥 🚳 🕂 🏐 🌴

#### Food, Lumber, Medicinal, Product, Ceremonial

Produces latex. The ripe fruit can be eaten raw and is rubbed between the palms of the hand as bruising the flesh eliminates astringency. It is also served in 'rujak', a fruit salad with a spicy sauce, pickled, or sweetened with sugar to make jam or confectionaries. The young leaves are eaten raw in side dishes as a vegetable. Used for magic. Immature fruit is used to prepare traditional medicine against diarrhea and dysentery. The juice of the leaves is applied to inflamed eye-lids. The wood is hard and strong and used to make household utensils such as pestles and furniture.

#### FARM SERVICES

Unknown

#### **BIODIVERSITY BENEFITS**

✓ YES Insects are attracted by the flower

Last Updated: February 17, 2021

Image: http://specimens.kew.org/herbarium/K000591189 The Herbarium Catalogue, Royal Botanic Gardens, Kew. Published on the Internet http://www.kew.org/herbcat [accessed on 10 12 2020]'.