

TREE SPECIES (SCIENTIFIC NAME)

Areca catechu

TREE FAMILY

ARECACEAE

AVERAGE LEAF SIZE (CM) 52.5cm × 5cm Length Width

ELEVATIONAL RANGE (M)

0-900м



ENGLISH COMMON NAME

Betelnut palm

INDONESIAN COMMON NAME

Pinang, Jambe (Javanese, Sundanese), Pua (Lesser Sunda Islands, the Moluccas)

DISTRIBUTION



NATIVE TO

Region: Asia Philippines

EXOTIC IN

SE Asia: Cambodia, Laos, Malaysia, Papua New Guinea, Singapore, Thailand, Vietnam

Indonesia: Borneo, Java, Lesser Sunda Islands, Maluku Islands, New Guinea, Sulawesi, Sumatra

COFFEE AGROFORESTRY INFORMATION

COFFEE SYSTEM ROBUSTA () ARABICA	COFFEE IMPACT Unknown
REGENERATION	CULTIVATION

PREVALENCE

🧚 COMMON IN COFFEE AGROFORESTRY

TREE BENEFITS AND USES

FARMER USES



Food, Medicinal, Ornamental, Product, Ceremonial

The betel nut is chewed as mild stimulant, contains a large quantity of tannin and oils, and can be sold at market. The nut is applied externally to ulcers, sores, wounds, swellings and skin diseases, and it is used as an antiparasitic for humans and animals, to stimulate menstrual discharge, as a cure for diarrhea, urinary disorders, edema (swelling caused by fluid retention), and low back pain. The fragrant flowers are used in weddings and funerals, its wood used for fencing, and the tree used as a garden accessory.

TREE MANAGEMENT

Propagated exclusively by seed but can collect seedlings from fruits dropped by bats and squirrels and raise them into mature palms. Sow mature seeds as whole fruits directly after harvesting or after a few days of drying in semishade. Germinates after about 90 days. Select seeds from tree that bears fruit early and regularly, has a large number of leaves on the crown, and has high fruit set. Select fruits that are fully mature and float vertically in water with the top end upward. Seedlings require shade. At 12-24 months plants with five or more leaves can be planted with a ball of earth adhering to the roots. Plant at a depth of about 90 cm in well-drained soils. Requires regular weeding and mulching. Irrigation used in some drier areas and usually cultivated around homes and along upland farm borders. It requires a high, well-distributed annual rainfall of 1500-5000 mm.

FARM SERVICES

Unknown

BIODIVERSITY BENEFITS

Unknown

Last Updated: February 17, 2021

Image: Himesh Dilruwan Jayasinghe© Digital Image © Board of Trustees, RBG Kew http://creativecommons.org/licenses/by/3.0/