



TREE SPECIES (SCIENTIFIC NAME)

# *Shorea platyclados*

INDONESIAN COMMON NAME

Meranti bukit, Banio (Sumatra), Ketir (South Kalimantan), Meranti cingham (Eastern Sumatra)

TREE FAMILY

**DIPTEROCARPACEAE**

AVERAGE LEAF SIZE (CM)

**7.5cm × 2.5cm**  
Length Width

DISTRIBUTION



**NATIVE TO INDONESIA**

ELEVATIONAL RANGE (M)

**700–1350m**

TREE HEIGHT

**LARGE (> 35M)**



NATIVE TO

**Region:** Asia

Malaysia, Singapore, Thailand

**Indonesia:** Borneo, Sumatra

## COFFEE AGROFORESTRY INFORMATION

COFFEE SYSTEM



**ROBUSTA**



**ARABICA**

COFFEE IMPACT

Unknown

TREE MANAGEMENT

Propagated by seed. Sow seeds as soon as possible and recommended to soak the seed for 12 hours prior to sowing. When seedlings are 5 - 6cm tall (~2 weeks after germination) pot into individual 15 x 23 cm containers with good drainage holes. Use mixture of forest soil and sand (3:1 ratio) as potting medium in order to introduce the appropriate mycorrhiza to the roots. Place seedlings in 50 - 60% sunlight and water twice daily. Seedlings can be planted when 30 - 40cm tall and should be hardened off in full sunlight for one month prior to planting

REGENERATION

Unknown

CULTIVATION



**PLANTED**

PREVALENCE

Unknown

## TREE BENEFITS AND USES

FARMER USES



**Fuelwood, Lumber**

Wood is used for making bridges, wharves, beams, and for applications where great strength is required.

FARM SERVICES

Unknown

BIODIVERSITY BENEFITS

Unknown

Last Updated: February 4, 2021

Image: <http://specimens.kew.org/herbarium/K000700664> The Herbarium Catalogue, Royal Botanic Gardens, Kew. Published on the Internet <http://www.kew.org/herbcat> [accessed on 10 12 2020].